

LABEL: MAIN PANEL – FarmVet CYPER POUR-ON

FOR EXTERNAL ANIMAL USE ONLY

SLEGS VIR UITWENDIGE DIERGEBRUIK

FarmVet CYPER POUR-ON

Reg. No. G3985 Act/Wet 36/1947

Pour-on for the control of ticks as well as nuisance, biting and Tsetse flies on cattle.
Opgietmiddel vir die beheer van bosluise asook las-, bytende- en Tsetsevlieë op beeste.

Spreads quickly and gives effective cover.
Vinnige verspreiding en effektiewe dekking.

CAUTION • VERSIGTIG

Contains/Bevat:

Alphacypermethrin/Alfasipermetrien 2,25 % m/v
Piperonyl butoxide/Piperonilbutoksied 3,00 % m/v

100 mℓ, 200 mℓ, 250 mℓ, 500 mℓ, 1 ℓ, 5 ℓ, 10 ℓ, 20 ℓ

Registration holder/Registrasiehouer:

FARMVET ANIMAL HEALTH (PTY) LIMITED

Co./Mpy. Reg. No. 2011/006806/07

P O Box/Posbus 10413, ASTON MANOR 1630, Gauteng, R.S.A.

Tel/Fax: 011 396 2233

Lot No.:

Expires/Verval:

AMENDMENT TO LABEL (G3985)

1. Trade name change
 2. Registration holder: Company name change
- Submitted: April 2013 (1st Draft)

Page 1 of 3

STORAGE INSTRUCTIONS

Store in original container, firmly closed and protected from light, below 25°C
Keep away from food and feedstuffs.

WARNINGS

- WITHDRAWAL PERIODS:
Meat – 7 days
Milk – 48 hours
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- Wear protective clothing – waterproof apron and gloves.
- When using do not eat, drink or smoke.
- Wash hands and contaminated skin after use.
- Remove and wash contaminated clothing.
- Empty containers should be destroyed.
- If product is splashed on the eyes or face, wash with plenty of water.
- If swallowed rinse the mouth with water.

In both cases consult a doctor and make this label available.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED
WHERE POSSIBLE APPLY TO DRY ANIMALS.

Application:

- **General tick control**

1 m□/10 kg body mass:

5 – 10 m□ from ear base to ear base,

5 m□ to tail head,

remainder along the back from shoulders to rump.

- **Spot treatment**

Ticks

To control a flush of brown ear ticks:

5 – 10 m□ from ear base to ear base.

Tail: 5 m□ to the tail head.

BERGINGSAAWYSINGS

Berg in oorspronklike houer, dig geseël en beskerm teen lig, benede 25°C
Hou weg van voedsel en voere.

WAARSKUWINGS

- ONTREKKINGSPERIODES:
Vleis – 7 dae
Melk – 48 ure
- Hou buite bereik van kinders, oningeligte persone en diere.
- Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

VOORSORGMAATREËLS

- Dra beskermende klere – waterdigde voorskoot en handskoene.
- Moenie eet, drink of rook tydens hantering nie.
- Was hande en besoedelde vel na gebruik.
- Verwyder en was besoedelde klere.
- Leë houers moet vernietig word.
- As produk in aanraking met oë of gesig kom, was met baie water.
- As ingesluk, spoel mond met water.

In albei gevalle raadpleeg 'n dokter en maak hierdie etiket beskikbaar.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI
WAAR MOONTLIK DIEN TOE OP DROË DIERE.

Aanwending:

- **Algemene bosluisbeheer**

1 m□/10 kg liggaamsmassa:

5 – 10 m□ van oor tot oor,

5 m□ op stertwortel,

res op die ruglyn van skouers tot kruis.

- **Kolbehandeling**

Bosluse

Om 'n oorvloed van bruinoorbosluse te beheer:

5 – 10 m□ van oor tot oor.

Stert: 5 m□ op stertwortel.

- **Nuisance and biting flies**

10 m□ along the back for animals over 12 months of age, and

5 m□ for animals under 1 year of age.

Repeat every 4 – 8 weeks.

Herd treatment rather than treatment of individuals is advised.

- **Tsetse fly control**

1 m□/10 kg body mass along the back from shoulders to rump.

Repeat every 4 – 8 weeks.

Herd treatment rather than treatment of individuals is advised.

- **Las- en bytende vlieë**

10 m□ op die ruglyn vir diere ouer as 12 maande, en

5 m□ vir diere jonger as 1 jaar.

Herhaal elke 4 – 8 weke.

Behandeling van die hele kudde eerder as enkele diere word aanbeveel.

- **Tsetsevlieg beheer**

1 m□/10 kg liggaamsmassa op die ruglyn van skouers tot kruis.

Herhaal elke 4 – 8 weke

Dit word aanbeveel dat die hele kudde eerder as enkele diere behandel word.