

LABEL MAIN PANEL – FarmVet DELTA POUR-ON

FOR EXTERNAL ANIMAL USE ONLY

SLEGS VIR UITWENDIGE DIERGEbruik

FarmVet DELTA POUR-ON

Reg. No. G4002 Act/Wet 36/1947

Pour-on for the control of ticks as well as nuisance, biting and Tsetse flies on cattle.
Opgietmiddel vir die beheer van bosluise asook las-, bytende- en Tsetsevlieë op beeste.

CAUTION • VERSIGTIG

Contains/Bevat:

Deltamethrin/Deltametrien 1 % m/v
Piperonyl butoxide/Piperonilbutoksied 3,00 % m/v

100 mℓ, 200 mℓ, 250 mℓ, 500 mℓ, 1 ℓ, 5 ℓ, 10 ℓ, 20 ℓ

Registration holder/Registrasiehouer:

FARMVET ANIMAL HEALTH (PTY) LIMITED

Co./Mpy. Reg. No. 2011/006806/07

P O Box/Posbus 10413, ASTON MANOR 1630, Gauteng, R.S.A.

Tel/Fax: 011 396 2233

Lot No.:

Expires/Verval:

AMENDMENT TO LABEL (G4002)

1. Trade name change
 2. Registration holder: Company name change
- Submitted: April 2013 (1st Draft)

Page 1 of 3

STORAGE INSTRUCTIONS

Store in original container, firmly closed and protected from light, below 25°C
Keep away from food and feedstuffs.

WARNINGS

- WITHDRAWAL PERIODS:
Meat – 14 days.
Milk – 48 hours.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- Wear protective clothing
 - Do not eat, drink or smoke while applying the product.
 - Wash hands and contaminated skin after use.
 - Remove and wash contaminated clothing.
 - Empty containers should be destroyed.
 - If product is splashed on the eyes or face, wash with plenty of water.
 - If swallowed rinse the mouth with water.
- In both cases consult a doctor and make this label available.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED
FOR BEST RESIDUAL EFFECT APPLY WHEN CATTLE ARE DRY.

BERGINGSAAWYSINGS

Berg in oorspronklike houer, dig geseël en beskerm teen lig, benede 25°C
Hou weg van voedsel en voere.

WAARSKUWINGS

- ONTTREKKINGSPERIODES:
Vleis – 14 dae.
Melk – 48 ure.
- Hou buite bereik van kinders, oningeligte persone en diere.
- Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

VOORSORGMAATREËLS

- Dra beskermende klere.
 - Moenie eet, drink of rook tydens hantering van die produk nie..
 - Was hande en besoedelde vel na gebruik.
 - Verwyder en was besoedelde klere.
 - Leë houers moet vernietig word.
 - As produk in aanraking met oë of gesig kom, was met baie water.
 - As ingesluk, spoel mond met water.
- In albei gevalle raadpleeg 'n dokter en maak hierdie etiket besikbaar.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI VIR DOELTREFFENDE NAWERKING WEND AAN OP DROË DIERE.

Application:

- **General tick control**

1 m□/10 kg body mass:

5 – 10 m□ from ear base to ear base,
5 m□ to tail head and the remainder along
the back from shoulders to rump.

Apply every 7 days during heavy challenge.

- **Spot treatment**

Ticks

Brown ear ticks: 10 m□ from ear base to ear
base.

Bont- and redlegged ticks: 5 m□ to the tail
head.

- **Nuisance and biting flies**

10 m□ between the shoulders for animals
over 12 months of age, and

5 m□ for animals under 1 year of age.

- **Tsetse fly control**

1 m□/10 kg body mass along the upper mid
line from shoulders to rump.

Apply monthly or at longer intervals
depending on challenge.

It is important that all animals in the area are
treated.

Aanwending:

- **Algemene bosluisbeheer**

1 m□/10 kg liggaamsmassa:

5 – 10 m□ van oor tot oor,
5 m□ op stertwortel, en die res op die rug
van skouers tot kruis.

Wend elke 7 dae aan gedurende swaar
uitdaging.

- **Kolbehandeling**

Bosluisse

Bruinoorbosluisse: 10 m□ van oor tot oor.

Bont- en Rooipoot-bosluisse: 5 m□ aan
stertwortel.

- **Las- en bytende vlieë**

10 m□ tussen die skouers vir diere ouer
as 12 maande; en

5 m□ vir diere jonger as 1 jaar.

- **Tsetsevlieg beheer**

1 m□/10 kg liggaamsmassa bokant die
middellyn van skouers tot kruis.

Wend maandeliks aan of met langer
tussenposes afhangend van uitdaging.

Dit is belangrik dat alle diere in die
omgewing behandel word.