

LABEL: FarmVet ECONO DOSE  
250 mℓ AND 500 mℓ (WRAP-AROUND LABEL)  
IMMEDIATE CONTAINER – MAIN PANEL

FOR ANIMAL USE ONLY

SELGS VIR DIEREGERBUIK

**FarmVet ECONO DOSE** †  
Reg. No. G3967 Act/Wet 36/1947

Roundworm remedy for cattle, sheep, goats and ostriches.  
Rondewurmmiddel vir beeste, skape, bokke en volstruise.

**CAUTION • VERSIGTIG**

**Contains/Bevat**

Levamisole hydrochloride/Levamisoolhidrochloried 25 % m/v

<b>250 mℓ</b>	<b>500 mℓ</b>
---------------	---------------

**Registration holder/Registrasiehouer:**

FARMVET ANIMAL HEALTH (PTY) LIMITED

Co./Mpy. Reg. No. 2011/006806/07

P O Box/Posbus 10413, ASTON MANOR 1630, Gauteng, R.S.A.

Tel/Fax: 011 396 2233

† Imidothiazole/Imidotiasool

Lot No.:

Expires/Verval:

AMENDMENT TO LABEL (G3967)

1. Trade name change
  2. Registration holder: Company name change
- Submitted: April 2013 (1<sup>st</sup> Draft)

Page 1 of 3

LABEL: FarmVet ECONO DOSE  
250 m $\square$  AND 500 m $\square$  (WRAP-AROUND LABEL)  
IMMEDIATE CONTAINER – SIDE/BACK PANELS

## WARNINGS

- **DILUTE WITH WATER BEFORE USE.**
- DO NOT SLAUGHTER ANIMALS FOR HUMAN CONSUMPTION WITHIN 7 DAYS OF LAST TREATMENT.
- No worm remedy, not even FarmVet ECONO DOSE, should be dosed to sheep and goats that have not been immunized against pulpy kidney (enterotoxaemia).
- In case of poisoning consult a doctor and make this information available.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder

## DIRECTIONS FOR USE: USE ONLY AS DIRECTED

DILUTE THE CONTENTS OF ONE 500 m $\square$  PACK WITH 4,5  $\square$  OF WATER.  
AFTER DILUTION USE WITHIN 30 DAYS.  
DOES NOT STAIN WOOL OR MOHAIR.

### Dosage

Sheep and goats: 3 m $\square$ /10 kg body mass  
Cattle: 15 m $\square$ /50 kg body mass  
Ostriches: 12 m $\square$ /10 kg body mass

## WAARSKUWINGS

- **VERDUN MET WATER VOOR GEBRUIK.**

- MOENIE DIERE BINNE 7 DAE NA LAASTE BEHANDELING VIR MENSLIKE VERBRUIK SLAG NIE.
- In die geval van skape en bokke, moet geen wurmmiddel, nie eens FarmVet ECONO DOSE, gegee word aan diere wat nie teen bloednier geïmmuniseer is nie.
- In geval van vergiftiging raadpleeg 'n dokter en maak hierdie inligting beskikbaar.
- Hou buite die bereik van kinders, oningeligte persone en diere.
- Alhoewel hierdie produk breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

## GEBRUIKSAANYSINGS: GEBRUIK SLEGS SOOS AANGEDUI

VERDUN DIE INHOUD VAN EEN 500 m $\square$  VERPAKKING MET 4,5  $\square$  WATER.  
NA VERDUNNING GEBRUIK BINNE 30 DAE.  
VLEK NIE WOL OF BOKHAAR NIE.

### Dosis

Skape en bokke: 3 m $\square$ /10 kg  
liggaamsmassa  
Beeste: 15 m $\square$ /50 kg  
liggaamsmassa  
Volstruise: 12 m $\square$ /10 kg  
liggaamsmassa

**Example for cattle, sheep, and goats**

**Voorbeeld vir beeste, skape en bokke**

Body mass/ Liggaamsmassa	Dose/Dosis	Number of doses in pack/Aantal dosisse in verpakking	
		250 ml	500 ml
10 kg	3 m□	833	1 666
20 kg	6 m□	416	833
30 kg	9 m□	333	555
40 kg	12 m□	166	416
50 kg	15 m□	75	333
100 kg	30 m□	37	166

**EFFICACY**

**DOELTREFFENDHEID**

Worm species/Wurmsoort	Sheep/goats Skape/bokke		Cattle/Beeste	
	Immature/ Onvolwasse	Adult/ Volwasse	Immature/ Onvolwasse	Adult/ Volwasse
Wireworm/Haarwurm	*	*	*	*
Brown stomachworm/Bruinmaagwurm	*	*		
Long-necked bankruptworm/Langnekbankrotwurm	*	*		
Large-mouthed bowelworm/Grootbekwurm	*	*		
Nodular worm/Knoppieswurm	*	*	**	*
Lungworm/Longwurm	**	*	**	*
Hookworm/Haakwurm	*	*		*
Bankruptwom/Bankrotwurm ( <i>Trichostrongylus</i> spp.)	*	*		
Bankruptworm/Bankrotwurm ( <i>Cooperia</i> spp.)			**	*
			Ostriches/Volstruise	
Ostrich wireworm/Volstruis haarwurm			**	*

\* Controls/Beheer =  $\geq$  90 % effective/doeltreffend

\*\* Aids in control/Hulpmiddel in beheer = 60 – 89 % effective/doeltreffend

This list contains the more important endoparasite species. This anthelmintic may also be effective against other species. For more information consult your veterinarian.

Hierdie lys bevat die meer belangrike endoparasitiese spesies. Dit kan ook effektief teen ander spesies gebruik word. Vir verdere besonderhede raadpleeg u veearts.