

DISPLAY CARTON - SACHET

FOR EXTERNAL ANIMAL USE ONLY

SLEGS VIR UITWENDIGE DIERGEbruik

FarmVet FLU-POUR-ON

Reg. No. G3981 Act 36/1947

READY-TO- USE POUR-ON FOR TICKS ON CATTLE, SHEEP, OSTRICHES AND GAME GEREED-VIR-GEbruik OPGIETMIDDEL VIR BOSLUISE OP BEESTE, SKAPE, VOLSTRUISE EN WILD

- Controls ticks on cattle, sheep, ostriches and game.
- Controls Tsetse fly worry on cattle.
- Controls feather mite on ostriches.
- Sterilises female ticks.
- Easy application.
- Residual action.
- Non-systemic action.
- The product is rainfast.
- Beheer bosluise op beeste, skape, volstruise en wild.
- Beheer Tsetsevlieg plaë by beeste.
- Beheer veermyte op volstruise.
- Steriliseer wyfie-bosluise.
- Maklike toediening.
- Nawerking.
- Nie-sistemiese aksie.
- Die produk is reënvas.

HARMFUL • SKADELIK

Contains

Flumethrin/Flumetriën 1 % m/v

20 ml, 40 ml

Registration holder/Registrasiehouer:

FARMVET ANIMAL HEALTH (PTY) LIMITED

Co./Mpy. Reg. No. 2011/006806/07

P O Box/Posbus 10413, ASTON MANOR 1630, Gauteng, R.S.A.

Tel/Fax: 011 396 2233

Date of Manuf./Datum Verv.:

Lot No.:

Expiry/Verval:

AMENDMENT TO LABEL (G3981)

1. Trade name change
 2. Registration holder: Company name change
- Submitted: April 2013 (1st Draft)

Page 1 of 9

SACHET

FOR EXTERNAL ANIMAL USE ONLY

SLEGS VIR UITWENDIGE DIEREGERBUIK

FarmVet FLU-POUR-ON

Reg. No. G3981 Act 36/1947

READY-TO- USE POUR-ON FOR TICKS ON CATTLE, SHEEP, OSTRICHES AND GAME GEREED-VIR-GERBUIK OPGIETMIDDEL VIR BOSLUISE OP BEESTE, SKAPE, VOLSTRUISE EN WILD

- Controls ticks on cattle, sheep, ostriches and game.
- Controls Tsetse fly worry on cattle.
- Controls feather mite on ostriches.
- Sterilises female ticks.
- Easy application.
- Residual action.
- Non-systemic action.
- The product is rainfast.
- Beheer bosluise op beeste, skape, volstruise en wild.
- Beheer Tsetsevlieg plaie by beeste.
- Beheer veermyte op volstruise.
- Steriliseer wyfie-bosluise.
- Maklike toediening.
- Nawerking.
- Nie-sistemiese aksie.
- Die produk is reënvas.

HARMFUL • SKADELIK

Contains

Flumethrin/Flumetriën 1 % m/v

20 ml, 40 ml

FOR FULL INSTRUCTIONS SEE PACKAGE INSERT
VIR VOLLE BESONDERHEDE SIEN INGESLOTE VOUBILJET

Registration holder/Registrasiehouer:

FARMVET ANIMAL HEALTH (PTY) LIMITED

Co./Mpy. Reg. No. 2011/006806/07

P O Box/Posbus 10413, ASTON MANOR 1630, Gauteng, R.S.A.

Tel/Fax: 011 396 2233

Lot No.:

Expiry/Verval:

PACKAGE INSERT - SACHET

FOR EXTERNAL ANIMAL USE ONLY

FarmVet FLU-POUR-ON

Reg. No. G3981 Act 36/1947

INDICATIONS

READY-TO- USE POUR-ON FOR TICKS ON CATTLE, SHEEP, OSTRICHES AND GAME

- Controls ticks on cattle, sheep, ostriches and game.
- Controls Tsetse fly worry on cattle.
- Controls feather mite on ostriches.
- Sterilises female ticks.
- Easy application.
- Residual action.
- Non-systemic action.
- The product is rainfast.

COMPOSITION

Flumethrin 1 % m/v

STORAGE INSTRUCTIONS

Protect from extremes of heat, cold and from dampness.

Store away from food and feed.

Tightly reseal the container if all the contents have not been used.

WARNINGS

- WITHDRAWAL PERIOD FOR MILK: NO WITHDRAWAL.
- WITHDRAWAL PERIOD FOR MEAT: 7 DAYS.
- Toxic to bees and fish.
- Although the colour of the product may vary, it will not influence the efficacy or residual period thereof.
- Skin reactions may occur in a small percentage of cases. If this happens, wash the affected animals with soap and water.
- Keep out of reach of uninformed persons, children and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- Do not eat, drink or smoke during or after handling this product, before washing hands and face thoroughly.
- Avoid contact with pour-on and wash immediately and thoroughly with soap and hot water after accidental contamination.
- Remove and wash all contaminated clothing.
- Destroy empty containers and do not re-use for any other purpose.
- Do not contaminate dams, rivers and other water sources.

AMENDMENT TO LABEL (G3981)

1. Trade name change
 2. Registration holder: Company name change
- Submitted: Aprii 2013 (1st Draft)

SLEGS VIR UITWENDIGE DIEREGERBUIK

FarmVet FLU-POUR-ON

Reg. No. G3981 Wet 36/1947

AANWYSINGS

GEREED-VIR-GEbruik OPGIETMIDDEL VIR BOSLUISE OP BEESTE, SKAPE, VOLSTRUISE EN WILD

- Beheer bosluise op beeste, skape, volstruise en wild.
- Beheer Tsetsevlieg plaë by beeste.
- Beheer veermyte op volstruise.
- Steriliseer wyfie-bosluise.
- Maklike toediening.
- Nawerking.
- Nie-sistemiese aksie.
- Die produk is reënvas.

SAMESTELLING

Flumetriën 1 % m/v

BERGINGSAAANWYSINGS

Beskerm teen uiterstes van hitte, koue en teen klammigheid.

Berg weg van voedsel en voer.

Indien inhoud nie opgebruik word nie, herseël houër deeglik.

WAARSKUWINGS

- ONTTREKKINGSPERIODE VIR MELK: GEEN.
- ONTTREKKINGSPERIODE VIR VLEIS: 7 DAE.
- Giftig vir bye en visse.
- Alhoewel die produk se kleur mag varieer, sal dit geen invloed hê op die doeltreffendheid of nawerking van die produk nie.
- Velreaksie mag in 'n klein persentasie van gevalle verwag word. Indien dit gebeur, was die geaffekteerde diere met seep en water.
- Hou buite bereik van oningeligte persone, kinders en diere.
- Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

VOORSORGMAATREËLS

- Moenie eet, drink of rook tydens of na aanwending van die middel alvorens die hande en gesig deeglik gewas is nie.
- Vermoedlik met opgietsmiddel en was onmiddellik en deeglik met seep en warm water na toevallige besmetting.
- Verwyder en was besoedelde klere.
- Vernietig leë houers en moenie vir enige ander doel gebruik nie.
- Moenie damme, riviere en ander waterbronne besoedel nie.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

- Treat at regular interval according to tick challenge.
- The longer the product is regularly applied, the more the tick population is reduced. It may then become possible to treat at extended intervals.

Dosage

The basic dosage is 1 m^l per 10 kg body mass (refer to dosage table below).

- **Cattle:** Apply the required dose in two lines, from the base of the skull along both sides of the spine up to the tail root. For Tsetse fly treat every 14 days or according to challenge.
- **Sheep:** Apply the required dose volume in 4 parts to the bare skin areas of the groin and axillae. For **prevention** of Karoo Paralysis, Coopers Redline has to be applied early in the season when the Karoo-Paralysis tick is first observed (either on the grazing or on the animals). Two to three treatments 7 to 10 days apart at this time should protect the sheep against Karoo-paralysis, depending on the severity of the tick challenge. For **treatment** of Karoo Paralysis (when the sheep are already recumbent), pour-on dip formulations are not recommended, as it takes a while for the product to distribute over the skin. Ensure immediate detachment of **all** ticks by spraying the belly and legs of affected animals with an amitraz-based product such as FarmVet AMITRAZ 20 % (G3981).
- **Game:** Apply the required dose volume with a suitable applicator. If self-medicating devices are used, it is important that the device only remains activated for as long as it takes to deliver one dosage to the animals in a particular paddock / camp. The device should not be used for continuous, low-level treatment.
- **Ostriches:** Apply the required dose volume along the backline.

Dosage table/Dosis tabel**Cattle and Game/Beeste en Wild**

Body mass/Liggaansmassa	Dosage/Dosis
Up to/Tot 100 kg	1 m ^l per 10 kg
100 kg	10 m ^l
200 kg	20 m ^l
300 kg	30 m ^l
400 kg	40 m ^l

Registration holder/Registrasiehouer:

FARMVET ANIMAL HEALTH (PTY) LIMITED

Co./Mpy. Reg. No. 2011/006806/07

P O Box/Posbus 10413, ASTON MANOR 1630, Gauteng, R.S.A.

Tel/Fax: 011 396 2233

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI

- Behandel teen gereelde tussenposes na gelang van die bosluisuitdaging.
- Hoe langer die produk gereeld gebruik word, hoe groter die daling in die bosluis bevolking. Dit mag dan moontlik wees om teen verlengde periodes te behandel.

Dosis

Die basiese dosis is 1 m^l/10 kg liggaansmassa.

- **Beeste:** Dien die vereiste dosis toe in twee strepe, vanaf die basis van die kop langs beide kante van die ruggraat tot op die stertwortel. Vir Tsetse vlieg, behandel met 14 dae tussenposes of na gelang van die vliegplaag.
- **Skape:** Dien die vereiste dosis in 4 dele toe aan die gedeeltes van die lieste en oksels waar die vel kaal is. Om Karoo-verlamming te **voorkom**, moet Coopers Redline aangewend word vroeg in die seisoen, wanneer die bosluis vir die eerste maal opgemerk word (op die weiding of op die diere). Twee of drie behandelings met 'n 7 tot 10 dae tussenpose op hierdie tyd behoort voldoende beskerming teen Karoo-verlamming te bied afhangend van die graad van bosluisbesmetting. Om Karoo-verlamming te **behandel** (wanneer die skape alreeds lê), word opgieter dip formulasies nie aanbeveel nie omrede die produk 'n tyd lank neem om oor die vel te versprei. Verseker onmiddellike loslating van **alle** bosluis deur die pense en bene van besmette diere te bespuit met 'n amitraz-gebaseerde produk soos FarmVet AMITRAZ 20 % (G3981).
- **Wild:** Dien die vereiste dosis met 'n gepaste toediener toe. Indien 'n selftoediener gebruik word, is dit belangrik dat die toestel slegs geaktiveer bly vir solank dit nodig is om die diere in 'n betrokke kamp eenmaal te behandel. Die toestel moet nie vir langdurige, laevlak behandeling gebruik word nie.
- **Volstruise:** Wend die vereiste dosis met die ruglyn langs aan.

Dosage table/Dosis tabel**Sheep and Ostriches/Skape en Volstruise**

Body mass/Liggaansmassa	Dose/Dosis
20 kg	2 m ^l
30 kg	3 m ^l
40 kg	4 m ^l
50 kg	5 m ^l
60 kg	6 m ^l

LABEL: FARMVET FLU-POUR-ON
DISPLAY CARTON – MAIN PANEL

FOR EXTERNAL ANIMAL USE ONLY

SLEGS VIR UITWENDIGE DIEREGERBUIK

FarmVet FLU-POUR-ON

Reg. No. G3981 Act 36/1947

**READY-TO- USE POUR-ON FOR TICKS ON CATTLE, SHEEP, OSTRICHES AND GAME
GEREED-VIR-GEbruik OPGIETMIDDEL VIR BOSLUISE OP BEESTE, SKAPE, VOLSTRUISE
EN WILD**

- Controls ticks on cattle, sheep, ostriches and game.
- Controls Tsetse fly worry on cattle.
- Controls feather mite on ostriches.
- Sterilises female ticks.
- Easy application.
- Residual action.
- Non-systemic action
- The product is rainfast.
- Beheer bosluise op beeste, skape, volstruise en wild.
- Beheer Tsetsevlieg plae by beeste.
- Beheer veermyte op volstruise.
- Steriliseer wyfie-bosluise.
- Maklike toediening.
- Nawerking.
- Nie-sistemiese aksie.
- Die produk is reënvas.

HARMFUL • SKADELIK

Contains

Flumethrin/Flumetrien 1 % m/v

200 ml (500 ml, 1 □, 5 l, 6 l, 20 l)

Registration holder/Registrasiehouer:

FARMVET ANIMAL HEALTH (PTY) LIMITED

Co./Mpy. Reg. No. 2011/006806/07

P O Box/Posbus 10413, ASTON MANOR 1630, Gauteng, R.S.A.

Tel/Fax: 011 396 2233

Lot No.:

Expiry/Verval:

AMENDMENT TO LABEL (G3981)

1. Trade name change
 2. Registration holder: Company name change
- Submitted: April 2013 (1st Draft)

Page 5 of 9

LABEL: FARMVET FLU-POUR-ON
DISPLAY CARTON – SIDE PANELS

STORAGE INSTRUCTIONS

Protect from extremes of heat, cold and from dampness.

Store away from food and feed.

Tightly reseal the container if all the contents have not been used.

WARNINGS

- WITHDRAWAL PERIOD FOR MILK: NO WITHDRAWAL.
- WITHDRAWAL PERIOD FOR MEAT: 7 DAYS.
- Toxic to bees and fish.
- Although the colour of the product may vary, it will not influence the efficacy or residual period thereof.
- Skin reactions may occur in a small percentage of cases. If this happens, wash the affected animals with soap and water.
- Keep out of reach of uninformed persons, children and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- Do not eat, drink or smoke during or after handling this product, before washing hands and face thoroughly.
- Avoid contact with pour-on and wash immediately and thoroughly with soap and hot water after accidental contamination.
- Remove and wash all contaminated clothing.
- Destroy empty containers and do not re-use for any other purpose.
- Do not contaminate dams, rivers and other water sources.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

- SHAKE WELL BEFORE USE.
- Treat at regular interval according to tick challenge.
- The longer the product is regularly applied, the more the tick population is reduced. It may then become possible to treat at extended intervals.

Dosage

The basic dosage is 1 mL per 10 kg body mass (refer to dosage table below).

- **Cattle:** Apply the required dose in two lines, from the base of the skull along both sides of the spine up to the tail root. For Tsetse fly treat every 14 days or according to challenge.
- **Sheep:** Apply the required dose volume in 4 parts to the bare skin areas of the groin and axillae. In the case of sheep infested with Karoo-paralysis tick (*Ixodes rubicundus*), VillVet FLU-POUR-ON has to be applied early in the season when these ticks are first observed (either on the grazing or on the animals). Two to three treatments 7 to 10 days apart at this time should protect the sheep against Karoo-paralysis, depending on the severity of the tick challenge.
- **Game:** Apply the required dose volume with a suitable applicator. If self-medicating devices are used, it is important that the device only remains activated for as long as it takes to deliver one dosage to the animals in a particular paddock / camp. The device should not be used for continuous, low-level treatment.

- **Ostriches:** Apply the required dose volume along the backline.

Dosage table: Cattle and Game.

Body mass	Dosage	No. of doses per ℓ
Up to 100 kg	1 m□ per 10 kg	
100 kg	10 m□	100
200 kg	20 m□	50
300 kg	30 m□	33
400 kg	40 m□	25
500 kg and over	50 m□	20

Dosage table: Sheep and Ostriches.

Body mass (kg)	Dose	No. of doses per ℓ
20	2 ml	500
30	3ml	333
40	4ml	250
50	5ml	200
60	6ml	166

Applicator

Squeeze bottle with dose meter. Squeeze required volume into dose meter, or use your drench gun.

BERGINGSAAWYSINGS

Beskerm teen uiterstes van hitte, koue en teen klammigheid.

Berg weg van voedsel en voer.

Indien inhoud nie opgebruik word nie, herseël houer deeglik.

WAARSKUWINGS

- ONTTREKKINGSPERIODE VIR MELK: GEEN.
- ONTTREKKINGSPERIODE VIR VLEIS: 7 DAE.
- Giftig vir bye en visse.
- Alhoewel die produk se kleur mag varieer, sal dit geen invloed hê op die doeltreffendheid of nawerking van die produk nie.
- Velreaksie mag in 'n klein persentasie van gevalle verwag word. Indien dit gebeur, was die geaffekteerde diere met seep en water.
- Hou buite bereik van oningeligte persone, kinders en diere.
- Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

VOORSORGMAATREËLS

- Moenie eet, drink of rook tydens of na aanwending van die middel alvorens die hande en gesig deeglik gewas is nie.
- Vermy aanraking met opgiemiddel en was onmiddellik en deeglik met seep en warm water na toevallige besmetting.
- Verwyder en was besoedelde klere.
- Vernietig leë houers en moenie vir enige ander doel gebruik nie.
- Moenie damme, riviere en ander waterbronne besoedel nie.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI

- SKUD GOED VOOR GEBRUIK.
- Behandel teen gereelde tussenposes na gelang van die bosluisuitdaging.
- Hoe langer die produk gereeld gebruik word, hoe groter die daling in die bosluis bevolking. Dit mag dan moontlik wees om teen verlengde periodes te behandel.

Dosis

Die basiese dosis is 1 ml/10 kg liggaamsmassa.

- **Beeste:** Dien die vereiste dosis toe in twee strepe, vanaf die basis van die kop langs beide kante van die ruggraat tot op die stertwortel. Vir Tsetse vlieg, behandel met 14 dae tussenposes of na gelang van die vliegplaag.
- **Skape:** Dien die vereiste dosis in 4 dele toe aan die gedeeltes van die lieste en oksels waar die vel kaal is. In die geval van skape wat aan besmetting met die Karoo-verlamings bosluis (*Ixodes rubicundus*) blootgestel word, moet FarmVet FLU-POUR-ON aangewend word vroeg in die seisoen, wanneer die bosluise vir die eerste maal opgemerk word (op die weiding of op die diere). Twee of drie behandelings met 'n 7 tot 10 dae tussenpose op hierdie tyd behoort voldoende beskerming teen Karooverlamming te bide afhangend van die graad van bosluisbesmetting.

- **Wild:** Dien die vereiste dosis met 'n gepaste toediener toe. Indien 'n selftoediener gebruik word, is dit belangrik dat die toestel slegs geaktiveer bly vir solank dit nodig is om die diere in 'n betrokke kamp eenmaal te behandel. Die toestel moet nie vir langdurige, laevlak behandeling gebruik word nie.
- **Volstruise:** Wend die vereiste dosis met die ruglyn langs aan.

Dosis tabel: Beeste en Wild.

Liggaamsmassa	Dosis	No. dosisse in 1 ℓ
Tot 100 kg	1 m□ per 10 kg	
100 kg	10 m□	100
200 kg	20 m□	50
300 kg	30 m□	33
400 kg	40 m□	25
500 kg	50 m□	20

Dosis tabel: Skape en Volstruise.

Liggaamsmassa (kg)	Dosis	No. dosisse in 1 ℓ
20	2ml	500
30	3ml	333
40	4ml	250
50	5ml	200
60	6ml	166

Toediener

Druk die vereiste volume uit die plastiese bottel in die maatbekertjie in, of gebruik 'n doseerspuit.