

OUTER CONTAINER: FarmVet OXYJECT

FOR ANIMAL USE ONLY

SLEGS VIR DIEREGERBUK

FarmVet OXYJECT

Reg. No. G3966 Act/Wet 36/1947

BROAD SPECTRUM AND LONG ACTING INJECTABLE ANTIBIOTIC BREE-SPEKTRUM EN LANG WERKENDE INSPUITBARE ANTIBIOTIKA

For use in cattle, sheep, goats and pigs.

For the treatment of conditions caused by, or associated with oxytetracycline susceptible organisms.

- **Cattle:** Treatment of tick-borne gallsickness (anaplasmosis), heartwater, pneumonia, footrot, joint-ill, navel-ill and pink eye (infectious bovine keratoconjunctivitis).
- **Sheep and goats:** Treatment of heartwater, pneumonia, footrot, joint-ill and navel-ill.
- **Pigs:** Treatment of pneumonia, joint-ill and navel-ill.

Vir gebruik in beeste, skape, bokke en varke.

Vir die behandeling van siektes veroorsaak, of geassosieer met oksitetrasiklien vatbare organismes.

- **Beeste:** Behandeling van bosluisoorge-draagde galsiekte (anaplasmose), hartwater, longontsteking, vrotpootjie, septiese gewrigsontsteking, naelstring-ontsteking en oogslymvliesontsteking (aansteeklike keratokonjunktivitis)
- **Skape en bokke:** Behandeling van hartwater, longontsteking, vrotpootjie, septiese gewrigsontsteking en naelstringontsteking.
- **Varke:** Behandeling van longontsteking, septiese gewrigsontsteking en naelstringontsteking.

STORAGE INSTRUCTIONS

Store at 5 – 25 °C in a dry place and out of direct sunlight.

BERGINGSAAWYSINGS

Berg tussen 5 – 25 °C in a droë plek weg van direkte sonlig.

Contains/Bevat:

Oxytetracycline/Oksitetrasiklien 20 g/100 ml

40ml, 100 ml, 250 ml, 500 ml

FOR FULL PARTICULARS REFER TO ENCLOSED PACKAGE INSERT
VIR VOLLE BESONDERHEDE SIEN INGESLOTE VOUBILJET

Registration holder/Registrasiehouer:

FARMVET ANIMAL HEALTH (PTY) LIMITED

Co./Mpy. Reg. No. 2011/006806/07

P O Box/Posbus 10413, ASTON MANOR 1630, Gauteng, R.S.A.

Tel/Fax: 011 396 2233

Lot No.:

Expires/Verval:

AMENDMENT TO LABEL (G3966)

1. Trade name change
 2. Registration holder: Company name change
- Submitted: April 2013 (1st Draft)

Page 1 of 5

IMMEDIATE CONTAINER: FarmVet OXYJECT

FOR ANIMAL USE ONLY

SLEGS VIR DIEREGERBUK

FarmVet OXYJECT

Reg. No. G3966 Act/Wet 36/1947

BROAD SPECTRUM AND LONG ACTING INJECTABLE ANTIBIOTIC BREE-SPEKTRUM EN LANG WERKENDE INSPUITBARE ANTIBIOTIKA

For use in cattle, sheep, goats and pigs.

For the treatment of conditions caused by, or associated with oxytetracycline susceptible organisms.

- **Cattle:** Treatment of tick-borne gallsickness (anaplasmosis), heartwater, pneumonia, footrot, joint-ill, navel-ill and pink eye (infectious bovine keratoconjunctivitis).
- **Sheep and goats:** Treatment of heartwater, pneumonia, footrot, joint-ill and navel-ill.
- **Pigs:** Treatment of pneumonia, joint-ill and navel-ill.

Vir gebruik in beeste, skape, bokke en varke.

Vir die behandeling van siektes veroorsaak, of geassosieer met oksitetrasiklien vatbare organismes.

- **Beeste:** Behandeling van bosluisoorge-draagde galsiekte (anaplasmose), hartwater, longontsteking, vrotpootjie, septiese gewrigsontsteking, naelstring-ontsteking en oogslymvliesontsteking (aansteeklike keratokonjunktivitis)
- **Skape en bokke:** Behandeling van hartwater, longontsteking, vrotpootjie, septiese gewrigsontsteking en naelstringontsteking.
- **Varke:** Behandeling van longontsteking, septiese gewrigsontsteking en naelstringontsteking.

STORAGE INSTRUCTIONS

Store at 5 – 25 °C in a dry place and out of direct sunlight.

BERGINGSAAWYSINGS

Berg tussen 5 – 25 °C in a droë plek weg van direkte sonlig.

Contains/Bevat:

Oxytetracycline/Oksitetrasiklien 20 g/100 ml

40ml, 100 ml, 250 ml, 500 ml

FOR FULL PARTICULARS REFER TO ENCLOSED PACKAGE INSERT
VIR VOLLE BESONDERHEDE SIEN INGESLOTE VOUBILJET

Registration holder/Registrasiehouer:

FARMVET ANIMAL HEALTH (PTY) LIMITED

Co./Mpy. Reg. No. 2011/006806/07

P O Box/Posbus 10413, ASTON MANOR 1630, Gauteng, R.S.A.

Tel/Fax: 011 396 2233

Lot No.:

Expires/Verval:

AMENDMENT TO LABEL (G3966)

1. Trade name change
 2. Registration holder: Company name change
- Submitted: April 2013 (1st Draft)

Page 2 of 5

PACKAGE INSERT: FarmVet OXYJECT

FOR ANIMAL USE ONLY

FarmVet OXYJECT

Reg. No. G3966 Act 36/1947

BROAD SPECTRUM AND LONG ACTING INJECTABLE ANTIBIOTIC

INDICATIONS

For use in cattle, sheep, goats and pigs.

For the treatment of conditions caused by, or associated with oxytetracycline susceptible organisms.

- **Cattle:** Treatment of tick-borne gallsickness (anaplasmosis), heartwater, pneumonia, footrot, joint-ill, navel-ill and pink eye (infectious bovine keratoconjunctivitis).
- **Sheep and goats:** Treatment of heartwater, pneumonia, footrot, joint-ill and navel-ill.
- **Pigs:** Treatment of pneumonia, joint-ill and navel-ill.

COMPOSITION

Oxytetracycline 20 g/100 ml

STORAGE INSTRUCTIONS

Store at 5 – 25 °C in a dry place and out of direct sunlight.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

Administration

- **Cattle:** Not more than 20 ml per injection site.
- **Pigs:** Not more than 10 ml per injection site.
- **Sheep:** Not more than 5 ml per injection site.

Normally one injection is sufficient to treat disease. A second injection 3 – 5 days after the initial injection may be given if necessary. Consult your veterinarian.

AMENDMENT TO LABEL (G3966)

1. Trade name change
 2. Registration holder: Company name change
- Submitted: April 2013 (1st Draft)

SLEGS VIR DIEREGERBRUIK

FarmVet OXYJECT

Reg. No. G3966 Wet 36/1947

BREË-SPEKTRUM EN LANG WERKENDE INSPIITBARE ANTIBIOTIKA

AANWYSINGS

Vir gebruik in beeste, skape, bokke en varke.

Vir die behandeling van siektes veroorsaak, of geassosieer met oksitetrasiklien vatbare organismes.

- **Beeste:** Behandeling van bosluisoorge-draagde galsiekte (anaplasmosis), hartwater, longontsteking, vrotpootjie, septiese gewrigsontsteking, naelstring-ontsteking en oogslimvliesontsteking (aansteeklike keratokonjunktivitis)
- **Skape en bokke:** Behandeling van hartwater, longontsteking, vrotpootjie, septiese gewrigsontsteking en naelstringontsteking.
- **Varke:** Behandeling van longontsteking, septiese gewrigsontsteking en naelstringontsteking.

SAMESTELLING

Oksitetrasiklien 20 g/100 ml

BERGINGSAAWYSINGS

Berg tussen 5 – 25 °C in a droë plek weg van direkte sonlig.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI

Toediening

- **Beeste:** Nie meer as 20 ml per inspuitplek.
- **Varke:** Nie meer as 10 ml per inspuitplek.
- **Skape:** Nie meer as 5 ml per inspuitplek.

Normaalweg is een inspuiting voldoende om siektes te behandel. 'n Tweede inspuiting 3 – 5 dae na die eerste inspuiting mag gegee word indien nodig. Raadpleeg u veearts.

Page 3 of 5

Dosage

- **Cattle, sheep and goats**
1 ml/10 kg bodymass - deep intramuscular injection.
- **Piglets up to 2kg bodymass**
0,25 ml by - subcutaneous injection.
- **Piglets up to 20 kg bodymass**
1 ml/animal – subcutaneous injection.
- **Pigs more than 20 kg bodymass**
1 ml/10 kg bodymass - deep intramuscular injection.

WARNINGS

- DO NOT USE IN DOGS.
- DO NOT USE IN HORSES.
- TREATED ANIMALS MUST NOT BE SLAUGHTERED FOR HUMAN CONSUMPTION WITHIN 28 DAYS OF LAST TREATMENT.
- MILK FROM TREATED ANIMALS MUST NOT BE USED WITHIN 5 DAYS OF LAST TREATMENT.
- DO NOT ADMINISTER TO PIGLETS ON THE SAME DAY AS THAT THEY ARE INJECTED WITH IRON.
- Slight inflammation may occur at the injection site. This will disappear within a few days.
- Hypersensitive animals may have allergic or anaphylactic reactions. Should this occur suspend treatment and consult your veterinarian.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

Dosis

- **Beeste, skape en bokke**
1 ml/10 kg liggaamsmassa - diep binnespiere inspuiting.
- **Klein varkies tot 2 kg liggaamsmassa**
0,25 ml - onderhuidse inspuiting.
- **Klein varkies tot 20 kg liggaamsmassa**
1 ml/dier – onderhuidse inspuiting
- **Varke meer as 20 kg liggaamsmassa**
1 ml/10 kg liggaamsmassa - diep binnespiere inspuiting.

WAARSKUWINGS

- MOENIE IN HONDE GEBRUIK NIE.
- MOENIE IN PERDE GEBRUIK NIE.
- BEHANDELDE DIERE MOENIE BINNE 28 DAE NA LAASTE BEHANDELING VIR MENSLIKE VERBRUIK GESLAG WORD NIE.
- MELK VAN BEHANDELDE KOEIE MOENIE VIR MENSLIKE VERBRUIK BINNE 5 DAE NA BEHANDELING GEBRUIK WORD NIE.
- MOENIE FarmVet OXYJECT OP DIESELFDE DAG BY VARKIES SPUIT AS WAT HUL MET YSTER INGESPUIT IS NIE.
- Geringe inflammasie mag by die inspuitingsplek voorkom. Dit sal na 'n paar dae verdwyn.
- Hipersensitiewe diere mag allergiese- of anafilaktiese reaksies toon. In dié geval staak behandeling en raadpleeg u veearts.
- Hou buite bereik van kinders, oningeligte persone en diere.
- Alhoewel hierdie middel breedvoerig onder 'n wye verskydenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

N.B. For rapid recovery a prompt and accurate diagnosis is essential. It is, therefore, always advisable to consult your veterinarian for a correct diagnosis.

Note: As with all oxytetracycline solutions darkening may occur during use without affecting efficacy.

Registration holder:

FARMVET ANIMAL HEALTH (PTY) LIMITED
Co. Reg. No. 2011/006806/07
P O Box 10413, ASTON MANOR 1630,
Gauteng, R.S.A.
Tel/Fax: 011 396 2233

L.W. Vir vinnige herstel is 'n spoedige en akkurate diagnose noodsaaklik. Dit is dus altyd raadsaam om u veearts vir 'n korrekte diagnose te raadpleeg.

Nota: Soos met alle oksitetrasiklien oplossings mag verdonkering gedurende gebruik voorkom sonder om die doeltreffendheid te beïnvloed.

Registration holder/Registrasiehouer:

FARMVET ANIMAL HEALTH (PTY) LIMITED
Mpy. Reg. No. 2011/006806/07
Posbus 10413, ASTON MANOR 1630,
Gauteng, R.S.A.
Tel/Fax: 011 396 2233