

LABEL: CONTAINER LABEL

VILLAVET SUPER WORM DOSE: 200 ml, 250 ml, 500 ml (WRAP-AROUND LABEL) AND 1 l, 5 l and 10 l

FOR ANIMAL USE ONLY

SLEGS VIR DIEREGERBRUIK

FarmVet SUPER WORM DOSE ③ + ⑧ †

Reg. No. G3980 Act/Wet 36/1947

Roundworm and tapeworm remedy for sheep, goats, cattle and ostriches.
Rondewurm en lintwurm behandeling vir skape, bokke, beeste en volstruise.

STORE IN A COOL, DRY PLACE.
BERG IN 'N KOEL, DROË PLEK.

CAUTION • VERSIGTIG

Contains/Bevat

Levamisole/Levamisool HCl 37,5 mg/ml

Praziquantel/Prasikwantel 18,8 mg/m²

200 ml (250 ml, 500 ml, 1 l, 5 l, 10 l)

Registration holder/Registrasiehouer:

FARMVET ANIMAL HEALTH (PTY) LIMITED

Co./Mpy. Reg. No. 2011/006806/07

P O Box/Posbus 10413, ASTON MANOR 1630, Gauteng, R.S.A.

Tel/Fax: 011 396 2233

† Imidothiazole/Imidotiasool
Isoquinoline/Isokwinalien

Lot No.:

Expiry/Verval:

AMENDMENT TO LABEL (G3980)

1. Trade name change
 2. Registration holder: Company name change
- Submitted: April 2013 (1st Draft)

Page 1 of 3

WARNINGS

- DO NOT SLAUGHTER ANIMALS FOR HUMAN CONSUMPTION WITHIN 7 DAYS OF TREATMENT.
- No worm remedy should be dosed to sheep and goats that have not been immunised against pulpy kidney.
- When treated with levamisole, the odd animal may develop an allergic-type reaction of a transient nature which normally disappears within 24 hours. An antihistamine may be administered. Under no circumstances must cortisone preparations be used.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

- **Sheep and goats:** 2 m□/10 kg body mass.
- **Cattle:** 10 m□/50 kg body mass.
- **Ostriches:** 1 m□/1 kg body mass.

WAARSKUWINGS

- MOENIE DIERE BINNE 7 DAE VANAF BEHANDELING VIR MENSLIKE VERBRUIK SLAG NIE.
- Geen wurmmiddel moet gedosser word in skape en bokke wat nie teen bloednier geënt is nie.
- Behandeling met levamisool mag lei na geïsoleerde gevalle van allergiese-tipe reaksies, wat van tydelike aard is en normaalweg binne 24 uur verdwyn. 'n Antihistamien mag toegedien word. Onder geen omstandighede moet kortisoon preperate gebruik word nie.
- Hou buite die bereik van kinders, oningeligte persone en diere.
- Alhoewel hierdie produk breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANBEVEEL

- **Skape en bokke:** 2 m□/10 kg liggaamsmassa.
- **Beeste:** 10 m□/50 kg liggaamsmassa.
- **Volstruise:** 1 m□/1kg liggaamsmassa.

EFFICACY

DOELTREFFENDHEID

Worm species/Wurmsort	Sheep/goats Skape/bokke		Cattle/Beeste	
	Immature/ Onvolwasse	Adult/ Volwasse	Immature/ Onvolwasse	Adult/ Volwasse
Wireworm/Haarwurm	*	*	*	*
Brown stomachworm/Bruinmaagwurm		**		**
Long-neck bankruptworm/Langnekbankrotwurm	**	*		
Large-mouthed bowelworm/Grootbekwurm	*	*		
Nodular worm/Knoppieswurm	*	*	**	*
Lungworm/Longwurm	**	**		
Hookworm/Haakwurm	*	*		*
Bankruptwom/Bankrotwurm (<i>Trichostrongylus</i> spp.)	*	*		
Bankruptworm/Bankrotwurm (<i>Cooperia</i> spp.)			*	*
Milk tapeworm/Melkintwurm	CLASS/KLAS I			
Wireworm/Haarwurm (<i>Libostrongylus douglassi</i>) Hookworm/Haakwurm (<i>Codiostomum struthionis</i>) Tapeworm/Lintwurm (<i>Houttuynia struthionis</i>)			Ostriches/Volstruise	
			**	*
			**	*
			**	*

AMENDMENT TO LABEL (G3980)

1. Trade name change
 2. Registration holder: Company name change
- Submitted: April 2013 (1st Draft)

* Controls/Beheer = \geq 90 % effective/doeltreffend

** Aids in control/Hulpmiddel in beheer = 60 – 89 % effective/doeltreffend

Class1: 100 % effective in more than 80 % of treated animals.

Klas I: 100 % doeltreffend in meer as 80 % van behandelde diere.

This list contains the more important endoparasite species. This anthelmintic may also be effective against other species. For more information consult your veterinarian.

Hierdie lys bevat die meer belangrike endoparasitiese spesies. Dit kan ook effektief teen ander spesies gebruik word. Vir verdere besonderhede raadpleeg u veearts.