

FOR EXTERNAL ANIMAL USE ONLY

SLEGS VIR UITWENDIGE DIEREGERBUK

### FarmVet WOUND REMEDY

Reg. No. G3986 Act/Wet 36/1947

#### SCREW-WORM AND WOUND REMEDY

A multi-purpose remedy which effectively controls and prevents screw-worm/blowfly strike, promotes healing and kills ticks present at time of application.

For use on livestock, horses, wild life, dogs, poultry and birds.

#### SPYKERWURM- EN WOND MIDDEL

'n Veeldoelige middel vir doeltreffende beheer en voorkoming van spykerwurm/brommer-maaier aanvalle, help met genesing en dood bosluise teenwoordig tydens aanwending.

Vir gebruik op vee, perde, wild, honde, pluimvee en voëls.

SHAKE WELL BEFORE USE □ SKUD GOED VOOR GEBRUIK

### CAUTION ♦ VERSIGTIG

- STORE BETWEEN 19 – 25 °C.
- CONTENTS FLAMMABLE AND UNDER PRESSURE.
- DO NOT STORE OR USE NEAR HEAT OR OPEN FLAMES.
- DO NOT LEAVE IN DIRECT SUNLIGHT.
- DO NOT PUNCTURE OR INCINERATE CONTAINER.
- BERG TUSSEN 19 – 25 °C.
- INHOUD ONTVLAMBAAR EN ONDER DRUK.
- MOENIE NABY HITTE OF OOP FLAMME BERG OF GEBRUIK NIE.
- MOENIE IN DIREKTE SONLIG LOS NIE.
- MOENIE HOER PERFOREER OF VERBRAND NIE.

#### Contains/Bevat

	% m/m
Resins and oils/Gom en olie	99,39
Germicide/Kiemdoder	0,51
Deltamethrin/Deltametrien	0,10

**190 g (300 mℓ)**

#### Registration holder/Registrasiehouer:

FARMVET ANIMAL HEALTH (PTY) LIMITED

Co./Mpy. Reg. No. 2011/006806/07

P O Box/Posbus 10413, ASTON MANOR 1630, Gauteng, R.S.A.

Tel/Fax: 011 396 2233

Lot No.:

Expires/Verval:

## FarmVet WOUND REMEDY – 190 g (300 ml) AEROSOL CAN SIDE PANEL

### WARNINGS

- DO NOT APPLY TO CATS.
- APPLY SPARINGLY TO BIRDS.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

### PRECAUTIONS

- Avoid inhalation of aerosol spray.
- Wash contaminated skin with soap and water.
- In the case of eye contact rinse immediately with copious amounts of water.
- In case of poisoning call a doctor and make this information available.
- Do not contaminate water sources or the environment.
- Dispose of containers in a safe manner and do not re-use for any other purpose.
- Store away from food and feed.

### DIRECTIONS FOR USE – USE ONLY AS DIRECTED DO NOT APPLY TO DEEP PENETRATING WOUNDS.

#### • All wounds

Apply in, around and above wounds (to diffuse downwards), to prevent screw-worm/blowfly strike and reduce bacterial infection and promote healing. This is recommended following routine procedures such as dehorning, knife castration, Mules operation and tail docking.

#### • Screw-worm/blowfly strike treatment

Apply freely into, around and above infested wounds. The worms will emerge within a short period, fall to the ground and die. With small wounds one treatment ± every 4 days until healed. Deeper wounds with extensive damage will respond more quickly if crusts and debris are gently removed before application but this is not essential. Better results follow application to animals which are dry.

### WAARSKUWINGS

- MOENIE NIE OP KATTE GEBRUIK NIE.
- GEBRUIK SPAARSAAM OP VOËLS.
- Hou buite bereik van kinders, oningeligte persone en diere.
- Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

### VOORSORGMATREËLS

- Vermyn inaseming van aërosol sprei.
- Was besoedelde kleres met seep en water.
- In geval van oogkontak spoel onmiddelik met baie water.
- In geval van vergiftiging raadpleeg 'n dokter en maak hierdie inligting beskikbaar.
- Moenie waterbronne of die omgewing besoedel nie.
- Doen weg met houers op 'n veilige manier en moenie vir enige ander doel hergebruik nie.
- Berg weg van kos en voer.

### GEBRUIKSAANWYSINGS – GEBRUIK SLEGS SOOS AANGEDUI

MOENIE AAN DIEP DEURDRINGENDE WONDE AANWEND NIE.

#### • Alle wonde

Wend aan in, om en bokant wonde (vir afwaartse verspreiding), vir voorkoming van spykerwurm/brommer-maaier aanvalle, vermindering van bakteriese infeksie en help met genesing. Bogenoemde word aanbeveel na prosedures soos onthoring, kastrasie, Mules operasie en stert afsit.

#### • Behandeling vir spykerwurm/brommer-maaier aanvalle

Gebruik ruimskoots in, om en bokant besmette wonde. Die wurms sal na 'n kort ruk verskyn, afval en vrek. Vir klein wonde, een behandeling ± elke 4 dae totdat dit genees is. Dieper wonde met baie beskadiging sal beter reageer as rowe en vuilgoed sagkens verwyder word voor aanwending, maar dit is nie noodsaaklik nie. Beter resultate sal verkry word as die middel op droë diere aangewend word.

FarmVet WOUND REMEDY

MAIN PANEL: 100 mℓ, 200 mℓ, 250 mℓ, 500 mℓ, (concertina label) and 1, 5 and 10 ℓ

FOR EXTERNAL ANIMAL USE ONLY

SLEGS VIR UITWENDIGE DIEREGERBUIK

**FarmVet WOUND REMEDY**

Reg. No. G3986 Act/Wet 36/1947

**SCREW-WORM AND WOUND REMEDY**

A multi-purpose remedy which effectively controls and prevents screw-worm/blowfly strike, promotes healing and kills ticks present at time of application.

For use on livestock, horses, wild life, dogs, poultry and birds.

**SPYKERWURM- EN WOND MIDDEL**

'n Veeldoelige middel vir doeltreffende beheer en voorkoming van spykerwurm/brommer-maaier aanvalle, help met genesing en dood bosluise teenwoordig tydens aanwending.

Vir gebruik op vee, perde, wild, honde, pluimvee en voëls.

**CAUTION ♦ VERSIGTIG**

**Contains/Bevat**

% m/m

Resins and oils/Gom en olie	99,39
Germicide/Kiemdoder	0,51
Deltamethrin/Deltametrien	0,10

**100 mℓ, 200 mℓ, 250 mℓ, 500 mℓ  
1 ℓ, 5 ℓ, 10 ℓ**

**Registration holder/Registrasiehouer:**

FARMVET ANIMAL HEALTH (PTY) LIMITED

Co./Mpy. Reg. No. 2011/006806/07

P O Box/Posbus 10413, ASTON MANOR 1630, Gauteng, R.S.A.

Tel/Fax: 011 396 2233

Lot No.:

Expires/Verval:

## FarmVet WOUND REMEDY

SIDE PANEL: 100 ml, 200 ml, 250 ml, 500 ml, (concertina label) and 1, 5 and 10 l

### WARNINGS

- DO NOT APPLY TO CATS.
- APPLY SPARINGLY TO BIRDS.
- Harmful if swallowed.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

### PRECAUTIONS

- Do not contaminate water sources or the environment.
- Dispose of containers in a safe manner and do not re-use for any other purpose.
- Store away from food and feed.
- Wash contaminated skin with soap and water.
- In the case of eye contact rinse immediately with copious amounts of water.
- In case of poisoning call a doctor and make this information available.

### DIRECTIONS FOR USE – USE ONLY AS DIRECTED DO NOT APPLY TO DEEP PENETRATING WOUNDS.

#### • All wounds

Apply in, around and above wounds (to diffuse downwards), to prevent screw-worm/blowfly strike and reduce bacterial infection and promote healing. This is recommended following routine procedures such as dehorning, knife castration, Mules operation and tail docking.

#### • Screw-worm/blowfly strike treatment

Apply freely into, around and above infested wounds. The worms will emerge within a short period, fall to the ground and die. With small wounds one treatment ± every 4 days until healed. Deeper wounds with extensive damage will respond more quickly if crusts and debris are gently removed before application but this is not essential. Better results follow application to animals which are dry.

### WAARSKUWINGS

- MOENIE NIE OP KATTE GEBRUIK NIE.
- GEBRUIK SPAARSAAM OP VOËLS.
- Skadelik indien ingesluk.
- Hou buite bereik van kinders, oningeligte persone en diere.
- Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

### VOORSORGMATREËLS

- Moenie waterbronne of die omgewing besoedel nie.
- Doen weg met houers op 'n veilige manier en moenie vir enige ander doel hergebruik nie.
- Berg weg van kos en voer.
- Was besoedelde klere met seep en water.
- In geval van oogkontak spoel onmiddelik met baie water.
- In geval van vergiftiging raadpleeg 'n dokter en maak hierdie inligting beskikbaar.

### GEBRUIKSAANWYSINGS – GEBRUIK SLEGS SOOS AANGEDUI MOENIE AAN DIEP DEURDRINGENDE WONDE AANWEND NIE.

#### • Alle wonde

Wend aan in, om en bokant wonde (vir afwaartse verspreiding), vir voorkoming van spykerwurm/brommer-maaier aanvalle, vermindering van bakteriese infeksie en help met genesing. Bogenoemde word aanbeveel na prosedures soos onthoring, kastrasie, Mules operasie en stert afsit.

#### • Behandeling vir spykerwurm/brommer-maaier aanvalle

Gebruik ruimskoots in, om en bokant besmette wonde. Die wurms sal na 'n kort ruk verskyn, afval en vrek. Vir klein wonde, een behandeling ± elke 4 dae totdat dit genees is. Dieper wonde met baie beskadiging sal beter reageer as rowe en vuilgoed sagkens verwyder word voor aanwending, maar dit is nie noodsaaklik nie. Beter resultate sal verkry word as die middel op droë diere aangewend word.